

# Child Maltreatment and Whole-person Health: Investigation of Psychosocial Buffers Using Structural Equation Modeling

Yanping Jiang, PhD, Emily A. Brinck, PhD, Sarah Ascienzo, PhD, Angela Groves, PhD, Kathy Trang, PhD, and Charlotte V. Farewell, PhD

**Objective:** The purpose of this study was to investigate psychosocial factors that may modify associations between child maltreatment and whole-person health (ie, depressive symptoms, chronic health conditions, psychological well-being, and inflammation).

**Methods:** Data were drawn from a subsample ( $N = 1255$ ) of Midlife in the United States study, a national longitudinal study of health and well-being. Structural equation modeling was applied to test hypotheses.

**Results:** Child maltreatment was associated with more depressive symptoms (standardized  $\beta = 0.29$ ,  $p < .001$ ), more chronic health conditions ( $\beta = 0.24$ ,  $p < .001$ ), poorer psychological well-being ( $\beta = -0.25$ ,  $p < .001$ ), and elevated C-reactive protein ( $\beta = 0.06$ ,  $p = .041$ ), but not with interleukin-6 (IL-6;  $\beta = 0.06$ ,  $p = .055$ ). There were significant interaction effects between child maltreatment and psychological resources on depressive symptoms (unstandardized  $b = -0.009$ ,  $SE = 0.005$ ,  $p = .045$ ) and IL-6 ( $b = -0.001$ ,  $SE = 0.000$ ,  $p = .042$ ). Specifically, there were stronger relationships between child maltreatment and more depressive symptoms and elevated IL-6 at low

levels of psychological resources than at high levels of psychological resources. There were no statistically significant interaction effects between child maltreatment and social resources on any health outcomes.

**Conclusions:** High levels of psychological resources significantly mitigate the long-term effects of child maltreatment on depressive symptoms and inflammation among middle-aged and older adults.

**Key Words:** biopsychosocial model, child maltreatment, whole person health, psychological resources, social resources, inflammation

**Abbreviations:** ACE = adverse childhood experiences, CRP = C-reactive protein, CTQ = Childhood Trauma Questionnaire, DMN = Default Mode Network, HPA = hypothalamic-pituitary-adrenal, IL-6 = interleukin-6, MIDUS = Midlife in the United States, SEM = structural equation modeling.

(*Biopsychosoc Sci Med* 2025;87:338–348)

Received for publication October 17, 2024; accepted March 12, 2025.

From the Department of Family Medicine and Community Health, Institute for Health, Health Care Policy and Aging Research, Rutgers, The State University of New Jersey, New Brunswick, New Jersey (Jiang); Wisconsin Center for Education Research, University of Wisconsin-Madison, Madison, Wisconsin (Brinck); School of Social Work, North Carolina State University, Raleigh, North Carolina (Ascienzo); Bronson School of Nursing, Western Michigan University, Kalamazoo, Michigan (Groves); Department of Epidemiology, Harvard T.H. Chan School of Public Health, Boston, Massachusetts (Trang); and Colorado School of Public Health, University of Colorado—Anschutz Medical Campus, Aurora, Colorado (Farewell).

Article Editor: David H. Chae

The analytical plan of this study was not pre-registered. The study protocol, materials, and de-identified data are available in a public archive: <http://www.icpsr.umich.edu/web/ICPSR/series/203>. The analytic code is not available in a public archive but will be made available upon request to Yanping Jiang, [yanping.jiang@ifh.rutgers.edu](mailto:yanping.jiang@ifh.rutgers.edu).

Address correspondence to Yanping Jiang, PhD, Department of Family Medicine and Community Health, Institute for Health, Health Care Policy and Aging Research, Rutgers, The State University of New Jersey, 112 Patterson Street, New Brunswick, NJ 08901. E-mail: [yanping.jiang@ifh.rutgers.edu](mailto:yanping.jiang@ifh.rutgers.edu)

Supplemental Digital Content is available for this article. Direct URL citations are provided in the HTML and PDF versions of this article on the journal's website, [journals.lww.com/bsam](http://journals.lww.com/bsam).

Copyright © 2025 Society for Biopsychosocial Science and Medicine  
DOI: 10.1097/PSY.0000000000001390

## INTRODUCTION

Well-being is a multifaceted construct that is directly related to the balance between an individual's resource pool and the psychological, social, and environmental contextual factors (eg, adverse stressful events) that may impact health.<sup>1</sup> However, the majority of social science research and practice continues to operationalize well-being as the absence of mental illness (eg, depression).<sup>2</sup> It is critical to redefine well-being beyond the absence of mental illness and explore multilevel mechanisms that foster overall health. This shift to a more salutogenic approach aligns with a focus on whole-person health, and consideration of how psychological (eg, optimism) and social (eg, social support) factors may buffer the experience of adverse stressful events, fostering mental and physical health and overall well-being.<sup>3,4</sup>

Child maltreatment, encompassing physical, emotional, and sexual abuse, as well as neglect,<sup>5,6</sup> is a critical issue with profound implications for whole-person health.<sup>7–9</sup> Globally, the prevalence of child maltreatment is alarming, with nearly 3 in 4 children aged 2 to 4 years experiencing physical punishment or psychological violence.<sup>10</sup> The health impacts of child maltreatment are immediate and long-lasting.<sup>11</sup> Findings from the Adverse

Childhood Experiences study revealed an individual's risk of developing mental and physical health conditions increased in a graded dose-response manner as the types of childhood adversity experienced increased.<sup>12</sup> Since the initial Adverse Childhood Experiences study, studies have continued to find that exposure to child maltreatment increases the risk for mental health problems, as well as increases the risk for myriad physical health diseases.<sup>13–17</sup>

According to the stress sensitization hypothesis, early life adversity, such as child maltreatment, can result in individuals having greater vulnerability to subsequent stressors, which in turn increases their risk for mental health problems such as depression in adulthood.<sup>18</sup> When applied to the biological embedding of early life adversity, the stress sensitization hypothesis suggests individuals who experience child maltreatment are at increased risk for subsequent health problems as a result of the structural and functional changes to the brain and biological systems caused by environmental stressors.<sup>19</sup> Specifically, child maltreatment experiences impact neurobiological processing pathways related to stress response, emotion processing, and self-referential processing.<sup>20</sup> Studies have identified alterations in the patterns of resting-state functional connectivity between these regions of the brain and between resting-state networks and revealed associations between psychopathology (eg, depression, post-traumatic stress disorder) and impaired functioning in these resting-state networks among those with child maltreatment histories.<sup>21,22</sup> In particular, growing evidence suggests that child maltreatment may negatively impact the Default Mode Network (DMN), which in turn leads to psychopathological symptoms and, particularly, those related to self-referential processing.<sup>23</sup> The DMN, sometimes referred to as the task-negative network, is involved in many high-level functions, including attention, self-related processes, and inhibitory control, and has an inverse relationship with the task-positive network.<sup>24</sup> Evidence suggests that child maltreatment experiences also impact the salience network, which processes attention, anticipation, or “alarm” and plays a crucial role in switching between the DMN and task-positive network.<sup>25</sup> Individuals with trauma histories and post-traumatic stress disorder have been found to have an overactive system that disproportionately assigns threats to stimuli.<sup>25</sup>

Studies suggest that changes in the stress response system, particularly the hypothalamic–pituitary–adrenal (HPA) axis, and inflammatory processes due to chronic stress exposure, drive the relationship between child maltreatment and long-term health outcomes.<sup>26</sup> Perceived stress activates the HPA axis, resulting in a cascade of responses, including the secretion of stress hormones, which increase inflammation. While initially adaptive, over time, this repeated or chronic activation of the stress response adversely affects HPA axis functioning and inflammatory processes.<sup>27</sup> In recent years, there has been a growing number of studies assessing associations between child maltreatment and inflammation, and recent systematic reviews and meta-analyses have helped to synthesize findings.<sup>28–30</sup> However, the authors caution that

methodological issues relating to the construct of child maltreatment, variations in analytic methods, and differences concerning consideration of confounding and mediating variables make it difficult to draw firm conclusions. These limitations notwithstanding, individuals with more severe child maltreatment histories tend to exhibit elevated inflammatory markers compared with those with less severe histories, and C-reactive protein (CRP) and interleukin-6 (IL-6) have emerged as markers most frequently assessed.<sup>28,30</sup> Inflammation plays a key role in many mental and physical health conditions and is considered both a consequence of child maltreatment and a risk factor for adverse health outcomes. Consequently, inflammation may be a key mechanism through which child maltreatment “gets under the skin” to affect long-term mental and physical health outcomes.<sup>31</sup> In addition, although there is a well-established association between individuals with major depressive disorder and elevated inflammation, inter-individual heterogeneity has been identified, leading some to assert elevated inflammatory markers may only be present for a subgroup of individuals with major depressive disorder, such as those who have experienced early life adversities.<sup>32</sup>

Despite increased understanding of the health-related consequences of child maltreatment, questions remain related to the pathways through which these experiences affect health, and how variations in the type of child maltreatment may differentially affect these processes and outcomes. While some studies have found maltreatment experiences involving threat (eg, physical abuse, sexual abuse, and psychological abuse), have worse impacts on health outcomes compared with experiences involving deprivation (eg, physical neglect and emotional neglect),<sup>33,34</sup> other studies have found significant relationships between health outcomes and experiences of threat or deprivation,<sup>35,36</sup> making additional inquiry necessary.

Importantly, the potential factors that might protect against adverse health outcomes associated with child maltreatment are not well understood yet. The stress-buffering hypothesis posits how psychosocial resources may mitigate the adverse effects of stress on individuals.<sup>37</sup> This hypothesis is crucial for developing interventions aimed at reducing stress-related health issues<sup>38,39</sup> The biopsychosocial model, which integrates biological, psychological, and social factors, provides a comprehensive framework for understanding health and illness, particularly in the context of stress.<sup>40</sup>

Psychological resources may mitigate maltreatment experienced in early life by contributing to positive mental dispositions and cognitive habits that are beneficial for well-being and health.<sup>41</sup> Self-esteem shapes individuals' health by influencing psychological, physical, and social well-being. Higher self-esteem is a psychological resource that helps individuals cope with adversity by regulating responses to the threat of stressors and thus supporting health.<sup>42</sup> A study by Dogan et al<sup>43</sup> determined that there were significant and positive relationships between both self-esteem and self-efficacy and psychological well-being.

Sense of control is another important psychological resource that has been shown to promote well-being, with individuals who have high self-control reporting more positive affect, life satisfaction, and happiness.<sup>44</sup> In addition, optimism is a frequently investigated psychological resource that promotes health and overall well-being. Higher optimism has been related prospectively to higher subjective well-being in times of adversity through greater engagement coping and less avoidance coping behaviors.<sup>45</sup> Moreover, a systematic review found that greater optimism was associated with increased engagement in health-promoting behaviors.<sup>46</sup> Collectively, these psychological resources may moderate associations between child maltreatment and whole-person health by promoting adaptive coping processes.

Greater social resources are also generally associated with more positive health outcomes,<sup>47,48</sup> and include consideration of 3 related but distinct constructs: social support, social strain, and social integration. Social support describes the perceived availability and quality of an individual's social support network, whereas social strain captures the perceived strain resulting from the demands of others (eg, partner, children, and friends). Social integration, in contrast, refers to the objective frequency of an individual's social interactions with high levels of contact (ie, integration) on one end of the spectrum and low levels of contact (ie, isolation) on the other. This is in line with prior literature, which has found that having larger social networks and higher quality of relationships—characterized by high support and low strain—are associated with better mental health<sup>49,50</sup> and physical health outcomes,<sup>47,48</sup> as well as well-being.<sup>51–53</sup> However, studies have tended to evaluate these social resources separately, and few have investigated relationships between the combined effects of social resources on the relationship between child maltreatment and health outcomes, overlooking the fact that these factors are largely overlapping and strongly correlated with each other.<sup>51</sup>

The stress-buffering hypothesis helps clarify the ways in which social resources may moderate associations between child maltreatment and health outcomes.<sup>37</sup> First, if individuals perceive they have others who will provide support in the aftermath of a stressful event, they may perceive the event as less stressful. Second, the availability of social support in the immediate aftermath of the event and longer term (eg, adulthood) may serve as a buffer between the event and the development of subsequent problems by providing individuals with resources (eg, emotional support) to more adaptively cope and potentially offset adverse impacts on the stress response system (ie, HPA axis) and inflammatory processes.<sup>15,26</sup> Safe and supportive relationships may be especially important for those who have experienced child maltreatment, as prior studies indicate maltreated youth are at elevated risk for exposure to other forms of violence.<sup>54,55</sup> In line with the stress-buffering hypothesis, social support has been found to buffer associations between stressful events and adverse mental health,<sup>53,56</sup> physical health,<sup>48,57</sup> and well-being outcomes.<sup>52</sup> To date, it remains unclear how collective

psychosocial resources may buffer against the deleterious effects of child maltreatment on mental health, physical health, well-being, and inflammation.

Social and economic consequences of child maltreatment are significant. Victims often face lower academic performance and reduced job prospects, contributing to an economic burden on society through increased healthcare costs and loss of productivity.<sup>9</sup> Addressing these issues requires a whole-person health approach, which integrates care for physical, mental, and psychological well-being.<sup>9</sup> In an attempt to fill this gap, the current study employed structural equation modeling (SEM) using nationally representative data from the Midlife in the United States (MIDUS 2) study to investigate psychological (ie, sense of control, self-esteem, and optimism) and social (ie, social integration, perceived social support, and low social strain) factors that may buffer associations between child maltreatment and whole-person health (ie, depressive symptoms, chronic health conditions, psychological well-being, and inflammation). We hypothesized that child treatment was associated with poorer whole-person health, and such an association was more evident among individuals with lower levels of psychological and social resources.

## METHODS

### Participants

A subsample of participants ( $N = 1255$ ) who participated in the biomarker project and completed the survey assessment at the second wave of the MIDUS 2 study 2004–2009 were included in our analyses.<sup>58</sup> The MIDUS study is an ongoing cohort study on the well-being and health of non-institutional adults with publicly available study protocol and data sets (accessed at <https://www.icpsr.umich.edu/web/ICPSR/series/203>). The biomarker project included a 24-hour stay at one of the 3 clinical research centers and the collection of fasting blood, saliva, and urine samples. The biomarker subsample reported higher levels of educational attainment and was less likely to smoke compared with the larger MIDUS 2 survey sample.<sup>59</sup> However, the biomarker subsample was comparable to the larger MIDUS 2 sample in terms of most key sociodemographic characteristics (eg, sex, age, race, marital status, income, and chronic conditions). Written consent was obtained from all participants, and the Institutional Review Boards at the University of Wisconsin–Madison, Georgetown University, and the University of California, Los Angeles approved the study.

### Measures

#### Child Maltreatment

Child maltreatment (the independent variable) was measured using the 28-item Childhood Trauma Questionnaire (CTQ) on a 5-point scale from 1 = never true to 5 = always true.<sup>60</sup> The CTQ is a validated measure of child maltreatment, consisting of the experience of emotional abuse, physical abuse, sexual abuse, emotional

neglect, and physical neglect. Of the 28 items, 3 validity items assessing minimization/denial were dropped from the analysis. Following previous MIDUS studies,<sup>61</sup> scores on each item were summed to create a composite, with higher scores reflecting higher levels of child maltreatment severity. In addition, 2 sum scores were created to reflect 2 distinct dimensions of child maltreatment: threat (ie, sum score of emotional, physical, and sexual abuse) and deprivation (ie, sum score of physical and emotional neglect).<sup>62</sup>

### Psychological Resources

The psychological resource latent factor (the moderator) was composed of 3 indicators: sense of control, self-esteem, and optimism. Sense of control was measured using a 12-item questionnaire<sup>63</sup> assessing personal mastery and perceived constraints on a 7-point scale from 1 = strongly agree to 7 = strongly disagree. A composite score was created by averaging responses on the 12 items, with higher scores reflecting higher levels of perceived control (Cronbach alpha = 0.86). Self-esteem was assessed using the 7-item version of the Rosenberg Self-Esteem Scale<sup>64</sup> on a 7-point scale from 1 = strongly agree to 7 = strongly disagree. A sum score was calculated, with higher scores reflecting higher levels of self-esteem (Cronbach alpha = 0.77). Optimism was measured using the 6-item life orientation test<sup>65</sup> on a 5-point scale from 1 = a lot agree to 5 = a lot disagree. A sum score was created, with higher scores reflecting higher levels of optimism (Cronbach alpha = 0.80).

### Social Resources

Three indicators were used to assess social resources (the moderator): social integration, social strain, and perceived social support. Following a similar approach adopted by previous MIDUS studies,<sup>66</sup> social integration was measured using 6 indicators: (1) married or not living alone, (2) weekly or more frequent contact with at least one family member, (3) weekly or more frequent contact with at least one friend, (4) weekly or more frequent contact with at least one neighborhood member, (5) monthly or more frequent attendance of church/temple activities, and (6) monthly or more frequent attendance of meetings of other social groups. Responses were summed to create a composite (range = 0 to 6), with higher scores reflecting higher levels of social integration. Social strain was measured using a 4-item scale asking participants' perception of strain (eg, making too many demands) from spouse/partner, family members, and friends, respectively.<sup>67</sup> Response options ranged from 1 = often to 4 = never. A mean score was calculated, with higher scores reflecting lower levels of social strain (Cronbach alpha = 0.82). Social support was measured using 4 items asking about their perceptions of support (eg, relying on for help) from spouse/partner, family members, and friends, respectively, on a 4-point scale from 1 = a lot and 4 = not at all.<sup>67</sup> Responses were reversely coded and averaged to create a composite, with higher scores reflecting higher levels of social support (Cronbach alpha = 0.84).

### Chronic Health Conditions

Chronic health conditions (dependent variable) were assessed by asking participants about the presence of each medical condition out of a list of 30 conditions (eg, cancer and diabetes) during the 12 months. A total count of the presence of each health condition was calculated, with higher scores reflecting more health conditions. Due to a skewed distribution, scores on chronic health conditions were log-transformed after adding a constant value of 1.

### Depressive Symptoms

Depressive symptoms (dependent variable) were assessed using the 20-item Center for Epidemiological Studies Depression Scale on a 4-point scale from 0 = rarely or none of the time to 3 = most or all of the time.<sup>68</sup> Responses were summed to create a composite score, with higher scores reflecting more depressive symptoms (Cronbach alpha = 0.89).

### Psychological Well-being

Following theoretical framework by Ryff,<sup>69</sup> psychological well-being (dependent variable) was indexed by 6 domains: autonomy, environmental mastery, personal growth, positive relations with others, purpose in life, and self-acceptance. Each domain was measured using 7 items, with response options ranging from 1 = strongly agree to 7 = strongly disagree. Responses on each set of items were summed to create a composite for each corresponding domain, with higher scores reflecting higher levels of psychological well-being subdomains (Cronbach alphas = 0.66 to 0.84).<sup>70</sup>

### Inflammation

Inflammation was indexed by CRP and IL-6 (dependent variables). CRP was determined using fasting blood through the BNII nephelometer (Dade Behring, inc.) by a particle-enhanced immunonephelometric assay.<sup>58</sup> For blood samples with undetectable CRP values, those samples were reassayed using a high-sensitivity kit (Meso Scale Diagnostics, #K151STG) by immunoelectrochemiluminescence. The inter-assay and intra-assay coefficients of variability ranged from 2.1% to 5.7%. IL-6 was determined using the Quantikine high-sensitivity enzyme-linked immunosorbent assay kit (R&D Systems). The inter-assay and intra-assay coefficients of variability, respectively, were 12.3% and 3.3%. Due to the skewed distribution, CRP and IL-6 were log-transformed for analyses.

### Covariates

Several key covariates were included in the analyses due to their potential effects on child maltreatment and health outcomes.<sup>71,72</sup> Covariates included sex (0 = male, 1 = female), race (0 = white, 1 = others), age, socioeconomic status, having physical activity for at least 20 minutes or more 3 times a week (0 = no, 1 = yes), alcohol use [0 = nonregular use (< 3 d/wk), 1 = regular use ( $\geq 3$  d/wk)], and smoking (0 = never smoked, 1 = current

or ever smoker). Socioeconomic status was operationalized by averaging z-scored highest levels of educational attainment (1 = no schooling/some grade schooling, 12 = doctoral and professional degree) and household-adjusted income. For models without the inclusion of social relationships, marital status (0 = others, 1 = married) was included as a covariate in the analysis.

## Statistical Analyses

Descriptive analyses were performed using SPSS 29.0. We used SEM to test our hypotheses in Mplus 8.11 using maximum likelihood with robust SE.<sup>73</sup> First, a measurement model (model 0) was performed to examine how well each observed indicator loaded on corresponding latent variables (ie, psychological well-being, psychological resources, and social resources). The model fit would be considered acceptable if the Comparative Fit Index is no smaller than 0.90 and the root mean square error of approximation is no larger than 0.08.<sup>74</sup> CRP and IL-6 were treated as 2 individual inflammatory markers in the analyses due to their distinct biological roles.<sup>75</sup> Child maltreatment severity, chronic health conditions, and depressive symptoms were also treated as observed variables in the SEM. Second, SEM (model 1) was performed by simultaneously testing the effect of child maltreatment on health outcomes (ie, depressive symptoms, psychological well-being, chronic health conditions, CRP, and IL-6) to minimize the inflation of type 2 errors. Finally, to test the moderating effect of psychological and social resources (model 2), psychological resources (or social resources) and the interaction term between child maltreatment and psychological resources (or social resources) were computed using the XWITH function and included in model 1. The moderating analyses were performed separately to minimize multicollinearity. Simple slope analyses were performed to further interpret significant moderation effects. Finally, we performed 2 sets of sensitivity analyses to examine whether the examined relationships might differ by different dimensions of child maltreatment. Specifically, we reran models 1 and 2 by treating threat and deprivation as independent variables in the analyses. Given the high correlation between threat and deprivation ( $r = 0.65$ ), we ran analyses by including threat and deprivation separately in the models. All models were performed by adjusting for covariates. Missing data were handled using the full information maximum likelihood, an efficient and robust approach to handling missing data in the SEM models.<sup>76</sup>

## RESULTS

Table 1 displays the characteristics of the sample. The mean age of the sample was  $54.52 \pm 11.72$  years. Over half of the sample (56.8%) were females, and the majority (77.2%) were white participants. Supplemental Digital Content, Table S1 (<http://links.lww.com/PSYMED/B88>), displays the bivariate correlations between observed variables of interest. Child maltreatment severity was negatively associated with each indicator of psychological and social resources, as well as indicators of

**TABLE 1.** Characteristics of the Study Sample

Variables	Mean	SD
Independent variable		
Child maltreatment severity	38.28	14.52
Threat	21.61	9.27
Deprivation	16.67	6.66
Moderators		
Social resources		
Social connectedness	4.34	1.23
Social strain	2.96	0.48
Social support	3.44	0.49
Psychological resources		
Optimism	23.62	4.80
Sense of control	5.59	0.99
Self-esteem	38.01	7.48
Dependent variables		
Chronic health conditions	2.47	2.48
CRP (ug/mL)	3.02	4.78
IL-6 (pg/mL)	3.04	3.04
Depressive symptoms	8.73	8.19
Psychological well-being		
Autonomy	37.34	6.69
Environmental mastery	38.24	7.70
Personal growth	39.44	6.76
Positive relations with others	40.50	7.23
Purpose in life	39.39	6.69
Self-acceptance	37.50	30.48
Covariates		
Regular physical activity (N, %)	960	76.5
Regular alcohol use (N, %)	237	18.9
Smoking (N, %)	597	47.6
Race (others; N, %)	286	22.8
Female (N, %)	713	56.8
Age	54.52	11.72
Household-adjusted income (\$)	36,380.4	31,928.7
Educational attainment	7.47	2.53

CRP = C-reactive protein; IL-6 = interleukin-6.

psychological well-being ( $p < .05$ ). Also, there were positive correlations between child maltreatment and chronic health conditions, CRP, IL-6, and depressive symptoms ( $p < .01$ ).

The measurement model (model 0) showed an acceptable model fit Satorra-Bentler  $\chi^2(51) = 624.92$ , Comparative Fit Index = 0.91, root mean square error of approximation = 0.08. Each observed indicator was significantly loaded on its corresponding latent variable ( $p < .001$ ), with factor loading values ranging from 0.43 to 0.86. Model 1, testing the main health effects of child maltreatment severity, showed that child maltreatment severity was associated with more depressive symptoms (standardized  $\beta = 0.29$ ,  $p < .001$ ), more chronic health conditions ( $\beta = 0.24$ ,  $p < .001$ ), poorer psychological well-being ( $\beta = -0.25$ ,  $p < .001$ ), and elevated CRP ( $\beta = 0.06$ ,  $p = .041$ ), but not with IL-6 ( $\beta = 0.06$ ,  $p = .055$ ), after adjusting for covariates (also see Supplemental Digital Content, Figure S1, <http://links.lww.com/PSYMED/B88>).

Moderation analyses (model 2) for psychological resources showed that there were significant interaction effects between child maltreatment severity and psychological resources on depressive symptoms (unstandardized  $b = -0.009$ , SE = 0.005,  $p = .045$ ) and IL-6 ( $b = -0.001$ ,

**TABLE 2.** The Moderating Effect of Psychological Resources

Variables	Psychological well-being		Depressive symptoms		Chronic health conditions		CRP		IL-6	
	B (SE)	p	B (SE)	p	B (SE)	p	B (SE)	p	B (SE)	p
CMS	-0.01 (0.01)	.095	0.09 (0.02)	<.001	0.01 (0.00)	<.001	0.01 (0.00)	.032	0.00 (0.00)	.27
PRs	0.91 (0.08)	<.001	-0.78 (0.18)	<.001	-0.05 (0.01)	.001	0.04 (0.03)	.15	0.02 (0.02)	.17
CMS × PRs	0.00 (0.00)	.19	-0.01 (0.01)	.045	0.00 (0.00)	.99	-0.00 (0.00)	.32	-0.00 (0.00)	.042
Regular physical activity	-0.26 (0.16)	.11	-0.76 (0.49)	.12	-0.02 (0.04)	.58	-0.44 (0.08)	<.001	-0.27 (0.05)	<.001
Regular alcohol use	0.02 (0.20)	.93	0.83 (0.56)	.14	-0.05 (0.05)	.27	-0.07 (0.08)	.44	-0.04 (0.05)	.47
Smoking	-0.14 (0.14)	.30	1.12 (0.10)	.005	0.09 (0.04)	.012	0.05 (0.07)	.42	0.07 (0.04)	.076
Race (others)	0.02 (0.18)	.90	1.57 (0.50)	.002	0.15 (0.05)	.003	0.23 (0.09)	.011	0.24 (0.06)	<.001
Female	0.39 (0.14)	.006	0.04 (0.40)	.92	0.18 (0.04)	<.001	0.33 (0.07)	<.001	0.06 (0.04)	.19
Age	0.02 (0.01)	.001	-0.03 (0.02)	.090	0.01 (0.00)	<.001	0.00 (0.00)	.51	0.01 (0.00)	<.001
Socioeconomic status	-0.02 (0.07)	.79	0.12 (0.25)	.64	0.01 (0.02)	.59	-0.11 (0.04)	.008	-0.06 (0.03)	.022
Marital status	0.18 (0.14)	.19	-1.39 (0.44)	.002	0.02 (0.04)	.72	0.06 (0.08)	.42	-0.07 (0.04)	.099

CMS = child maltreatment severity; CRP = C-reactive protein; IL-6 = interleukin-6; PRs = psychological resources.

SE = 0.000,  $p = .042$ ). Simple slope analyses showed that there was a stronger relationship between child maltreatment severity and depressive symptoms at low levels of psychological resources (SD = -1,  $b = 0.118$ , SE = 0.023,  $p < .001$ ) than at high levels of psychological resources (SD = 1,  $b = 0.054$ , SE = 0.022,  $p = .013$ ). Also, there was a significant relationship between child maltreatment severity and elevated IL-6 at low ( $b = 0.004$ , SE = 0.002,  $p = .014$ ) but not high ( $b = -0.001$ , SE = 0.002,  $p = .77$ ) levels of psychological resources. However, psychological resources did not moderate the relationship between child maltreatment severity and psychological well-being, chronic health conditions, or CRP (Table 2). There were no statistically significant interaction effects between child maltreatment severity and social resources on any health outcomes (Table 3).

**Sensitivity Analyses**

Similar to the results reported previously, threat and deprivation were associated with more depressive symptoms, more chronic health conditions, and poorer psychological well-being ( $p < .001$ ). Threat, but not deprivation, was associated with elevated CRP ( $\beta = 0.07$ ,

$p = .021$ ;  $\beta = 0.04$ ,  $p = .22$ ; respectively) and IL-6 ( $\beta = 0.06$ ,  $p = .039$ ;  $\beta = 0.04$ ,  $p = .19$ ; respectively). To examine whether threat and deprivation might have unique effects on these health outcomes, we included threat and deprivation simultaneously in model 1. Results showed that both threat and deprivation were associated with depressive symptoms ( $\beta = 0.09$ ,  $p = .030$ ;  $\beta = 0.23$ ,  $p < .001$ ; respectively). Threat, but not deprivation, was uniquely associated with chronic health conditions ( $\beta = 0.23$ ,  $p < .001$ ;  $\beta = 0.03$ ,  $p = .44$ ; respectively) and CRP ( $\beta = 0.07$ ,  $p = .050$ ;  $\beta = -0.01$ ,  $p = .75$ ; respectively). Deprivation, but not threat, was uniquely associated with psychological well-being ( $\beta = -0.28$ ,  $p < .001$ ;  $\beta = 0.00$ ,  $p > .99$ ; respectively). Neither threat nor deprivation was associated with IL-6 ( $\beta = 0.06$ ,  $p = .11$ ;  $\beta = -0.00$ ,  $p = .98$ ; respectively).

Sensitivity analyses for model 2 showed that psychological resources moderated the relationship between threat and depressive symptoms ( $b = -0.015$ , SE = 0.008,  $p = .041$ ) and the relationships between deprivation and psychological well-being ( $b = 0.007$ , SE = 0.003,  $p = .038$ ) and IL-6 ( $b = -0.002$ , SE = 0.001,  $p = .019$ ). Simple slope analyses showed that threat was associated

**TABLE 3.** The Moderating Effect of Social Resources

Variables	Psychological well-being		Depressive symptoms		Chronic health conditions		CRP		IL-6	
	B (SE)	p	B (SE)	p	B (SE)	p	B (SE)	p	B (SE)	p
CMS	-0.02 (0.01)	.091	0.10 (0.02)	<.001	0.01 (0.00)	<.001	0.01 (0.00)	.043	0.00 (0.00)	.21
SRs	3.58 (0.79)	<.001	-4.30 (1.59)	.007	-0.07 (0.11)	.52	0.24 (0.21)	.24	0.13 (0.12)	.28
CM × SRs	0.02 (0.02)	.49	-0.03 (0.05)	.50	-0.00 (0.00)	.44	-0.00 (0.00)	.43	-0.00 (0.00)	.17
Regular physical activity	0.08 (0.24)	.75	-1.18 (0.51)	.021	-0.05 (0.04)	.31	-0.43 (0.08)	<.001	-0.28 (0.05)	<.001
Regular alcohol use	-0.20 (0.27)	.47	1.15 (0.60)	.054	-0.04 (0.05)	.38	-0.07 (0.08)	.44	-0.03 (0.05)	.50
Smoking	0.04 (0.20)	.84	0.93 (0.43)	.029	0.08 (0.04)	.029	0.05 (0.07)	.42	0.08 (0.04)	.061
Race (others)	0.58 (0.28)	.041	1.28 (0.56)	.023	0.13 (0.05)	.012	0.23 (0.09)	.011	0.26 (0.05)	<.001
Female	0.21 (0.20)	.30	0.48 (0.44)	.27	0.18 (0.04)	<.001	0.31 (0.07)	<.001	0.06 (0.04)	.13
Age	0.04 (0.01)	<.001	-0.05 (0.02)	.002	0.01 (0.00)	<.001	0.00 (0.00)	.46	0.01 (0.00)	<.001
Socioeconomic status	0.67 (0.12)	<.001	-0.74 (0.26)	.004	-0.02 (0.02)	.39	-0.10 (0.04)	.016	-0.07 (0.03)	.009

CMS = child maltreatment severity; CRP = C-reactive protein; IL-6 = interleukin-6; SRs = social resources.

with depressive symptoms at low but not at high levels of psychological resources ( $b = 0.157$ ,  $SE = 0.037$ ,  $p < .001$ ;  $b = 0.049$ ,  $SE = 0.036$ ,  $p = .18$ ; respectively). Similarly, deprivation was associated with psychological well-being and IL-6 at low but not at high levels of psychological resources (psychological well-being:  $b = -0.060$ ,  $SE = 0.014$ ,  $p < .001$ ;  $b = -0.010$ ,  $SE = 0.020$ ,  $p = .62$ , respectively; IL-6:  $b = 0.009$ ,  $SE = 0.004$ ,  $p = .032$ ;  $b = -0.005$ ,  $SE = 0.005$ ,  $p = .28$ , respectively). Social resources did not interact with threat or deprivation to affect health outcomes ( $ps > .05$ ).

## DISCUSSION

This study applied the biopsychosocial model to investigate how psychosocial resources may buffer associations between child maltreatment severity and whole-person health in a nationally representative sample. Our findings suggest that child maltreatment was associated with more chronic health conditions, elevated inflammation, more depressive symptoms, and poorer psychological well-being. Moderation analyses indicated that though social resources did not significantly moderate these associations, psychological resources mitigated the impact of child maltreatment on depressive symptoms and inflammation (ie, IL-6).

Among this sample, child maltreatment was positively associated with elevated CRP in adulthood. This result is consistent with a meta-analysis by Baumeister et al<sup>28</sup>, showing a relationship between more severe child maltreatment histories and elevated CRP. Interestingly, our result did not support a significant relationship between child maltreatment severity and IL-6, though existing literature has shown elevated IL-6 among those who had experienced child maltreatment.<sup>77</sup> These results seem to suggest that compared with IL-6, CRP might be more sensitive to the long-term effects of early life experiences, including child maltreatment.<sup>30</sup>

We also found that child maltreatment was associated with more chronic health conditions and depressive symptoms, as well as worse psychological well-being. These findings align with prior research indicating that child maltreatment can lead to chronic emotional dysregulation and heightened vulnerability to physical and mental illness in adulthood.<sup>78–80</sup> In addition, our sensitivity analyses found that when threat and deprivation were included in the same model, threat was associated with CRP and chronic health conditions, and deprivation was associated with psychological well-being. These results, together with existing literature,<sup>62</sup> highlight the need to consider the distinct dimensions of child maltreatment and their potential impact on health.

Notably, our moderation analyses showed that psychological resources, including optimism, self-esteem, and sense of control, buffered the impact of child maltreatment severity on depressive symptoms and IL-6. Specifically, the associations between child maltreatment severity and depressive symptoms and IL-6 were stronger at low levels of psychological resources than at high levels

of psychological resources. This finding indicates that psychological resources may help mitigate the emotional and biological toll of child maltreatment,<sup>79</sup> although its protective effects may be limited in the context of more pervasive physical and psychological harm. These results align with a previous study that identified self-esteem as a psychological resource buffering the adverse effect of childhood victimization on mental health in adulthood.<sup>81</sup> Alternative research identified optimism as a psychological resource that mediated associations between child maltreatment and depression and highlighted the need for additional studies examining multilevel resources as moderators of mental health outcomes.<sup>82</sup>

A particularly novel finding of this study is the buffering effect of psychological resources on inflammation. While limited research has explored collective psychological resources in this context, mixed findings have emerged in the examination of sense of control as a moderator of trauma-related inflammation, reporting buffering effects<sup>21</sup> or null findings.<sup>83</sup> Our results corroborate the buffering effects of psychological resources, indicating that psychological resources play a protective role in buffering against the adverse effects of child maltreatment severity on inflammation, specifically by moderating the relationship with IL-6. This result suggests that individuals with greater psychological resources may experience reduced inflammatory responses linked to IL-6, potentially helping to mitigate the long-term health consequences of child maltreatment. However, the lack of moderation in the relationship between child maltreatment and CRP suggests that different inflammatory markers may be influenced by distinct mechanisms. Future research should examine the pathways through which psychological resources impact inflammatory responses and identify additional factors that may help build psychological resources in individuals with a history of child maltreatment.

Surprisingly, we found that social resources did not alleviate the negative consequences of child maltreatment on any health outcomes. Several factors, including the timing of social support, the embedding of the stress response, maladaptive coping mechanisms, and the participants' demographics, could explain these results. The timing of social support could be a crucial factor influencing our study results. For instance, social support and integration may be less effective when introduced later in life than continuous support from a younger age. If participants did not experience supportive relationships early in life to counterbalance the effects of maltreatment, it might be harder for them to benefit from such resources later on. This is supported by a recent study that underscores the importance of consistent early intervention, including social support and psychoeducational interventions, continuing into adulthood.<sup>84</sup> Notably, the ability to form supportive relationships can be impaired by child maltreatment, making individuals less likely to utilize available social resources effectively. Greater levels of child maltreatment may predict higher avoidance in close relationships.<sup>85</sup>

The biological embedding of stress responses due to child maltreatment might have impacted the results in the current study. Chronic exposure to stress due to child maltreatment can make it difficult for individuals to receive social support. This finding aligns with past research that found children from highly distressed households are less likely to develop the social and emotional skills necessary for forming supportive close relationships in adulthood.<sup>86</sup> Also, maladaptive coping mechanisms could have a significant influence on our study results. Participants with a history of child maltreatment may develop maladaptive coping mechanisms (eg, substance abuse, social withdrawal) that persist into adulthood and can undermine the benefits of social support. For example, participants with low social support and a diminished sense of belonging may be more susceptible to substance use.<sup>87</sup> Mental health disorders such as depression, which are often outcomes of child maltreatment, can interfere with the ability to form and maintain supportive social relationships, thus limiting the buffering effect of social support. Germane to this point, past research suggests a significant indirect effect of child maltreatment on depression, mediated by insecure attachment and perceived social support.<sup>85</sup>

### Limitations

A few limitations should be acknowledged when interpreting the findings of this study. First, although the CTQ is a commonly used, valid assessment for child maltreatment, reporting bias might occur in the retrospective self-report of child maltreatment experiences, resulting in overestimating or underestimating the health effects of child maltreatment. Existing studies have shown a moderate correlation between retrospective and prospective reports of child maltreatment and stronger relationships between retrospective reports of child maltreatment and mental health than prospective reports of child maltreatment.<sup>88,89</sup> Future studies are needed to validate our findings using prospective reports of child maltreatment. Also, this study solely relied on self-reports of child maltreatment. Contamination (eg, overreporting child maltreatment) may occur to obscure the reported relationship between child maltreatment severity and health.<sup>90</sup> Although Coelho et al<sup>29</sup> noted in their systematic review that the CTQ stands out as one of the more comprehensive measures of childhood trauma with strong psychometric properties, future studies might consider using multiple sources of reports on child maltreatment (eg, official case reports). Second, the cross-sectional nature of the data collection limits the ability to draw definitive conclusions about the causal relationships in this study. Third, differences in some characteristics (eg, educational attainment) between the biomarker subsample and the MIDUS survey sample might indicate the existence of a selection bias in our analytic sample that could affect the reported estimates. Fourth, social resource variables were not assessed during childhood when maltreatment occurred, but rather decades later, which precludes us from evaluating whether or how social resources

in childhood might have influenced long-term health outcomes. Fifth, we used depressive symptoms to represent mental health within our biopsychosocial model, and future studies should consider other mental health symptoms and disorders. Lastly, the analytic sample was predominantly white, which limits our ability to examine any meaningful racial differences in the relationships examined in this study.

### Implications

Despite these limitations, this study contributes additional information to understanding how child maltreatment affects health outcomes and provides novel insight into the protective role of psychological resources in mitigating the adverse effects of child maltreatment on whole-person health. Findings from this study advance current understanding of factors that may protect against the deleterious consequences of child maltreatment and hold implications for public health, particularly in the areas of prevention and intervention related to child maltreatment and whole-person health. The negative effects of child maltreatment on whole-person health in adulthood reported in this study highlight the importance of early intervention and support for individuals who experienced child maltreatment to reduce these adverse long-term health outcomes. For example, assessing the experience of child maltreatment should be considered as a routine screening for primary care, enabling early intervention and offering immediate support.<sup>78,91</sup> Moreover, this study shows that psychological resources (eg, self-esteem and sense of control) mitigated the adverse effects of child maltreatment on mental and immune function, suggesting that enhancing psychological resilience (eg, through cognitive behavioral therapy) could be a possible effective intervention for improving the mental and biological health of those with a history of child maltreatment. The finding that psychological resources buffered the effects of child maltreatment on depressive symptoms and inflammation, but not on overall psychological well-being or chronic health conditions suggests these resources may be more effective in reducing depressive symptoms and inflammatory responses as opposed to improving physical health or psychological well-being. In addition, the absence of significant interaction effects between social resources (eg, social support) and child maltreatment on health outcomes does not necessarily imply a less critical role of positive social relationships and social integration in fostering whole-person health than psychological resources. Our findings appear to suggest that social resources may not contribute to better health by protecting against the adverse health effects of child maltreatment. Instead, social resources might directly relate to better health outcomes, regardless of the presence of exposure to child maltreatment.

### CONCLUSION

In summary, this study provided important empirical evidence to support the protective effect of psychological resources in maintaining mental and biological health following exposure to child maltreatment. Findings

from this study suggest that interventions designed to enhance psychological resources might be a potentially effective strategy to promote health for middle-aged and older adults who experience child maltreatment.

*Source of Funding and Conflicts of Interest: Data collection for the present study was supported by the John D. and Catherine T. MacArthur Foundation Research Network on Successful Midlife Development, the National Institute on Aging (P01-AG020166), M01-RR023942 (Georgetown), M01-RR00865 (UCLA) from the General Clinical Research Centers Program, and 1UL1RR025011 (UW) from the Clinical and Translational Science Award (CTSA) program of the National Center for Research Resources, National Institutes of Health. The preparation of the manuscript was supported by the National Institute of Mental Health (R25MH136652). The authors report no conflicts of interest.*

## REFERENCES

- McNaught A, Knight A, McNaught A. Defining wellbeing. *Understanding wellbeing: an introduction for students and practitioners of health and social care*. Banbury: Scion Publishing Ltd; 2011: 7–23.
- Wadephul F, Glover L, Jomeen J. Conceptualising women's perinatal well-being: a systematic review of theoretical discussions. *Midwifery*. 2020;81:102598.
- Hefli R. The extended biopsychosocial model: a whole-person approach to psychosomatic medicine and psychiatry. *Psyche Geolof*. 2013;24:119–29.
- Frazier LD. The past, present, and future of the biopsychosocial model: A review of The Biopsychosocial Model of Health and Disease: new philosophical and scientific developments by Derek Bolton and Grant Gillett. *New Ideas Psychol*. 2020;57:100755.
- Massullo C, De Rossi E, Carbone GA, Imperatori C, Ardito RB, Adenzato M, et al. Child maltreatment, abuse, and neglect: an umbrella review of their prevalence and definitions. *Clin Neuro-psychiatry*. 2023;20:72.
- Vizard E, Gray J, Bentovim A. The impact of child maltreatment on the mental and physical health of child victims: a review of the evidence. *BJPsych Advances*. 2022;28:60–70.
- Fryers T, Brugha T. Childhood determinants of adult psychiatric disorder. *Clin Pract Epidemiol Ment Health*. 2013;9:1–50.
- Hughes K, Bellis MA, Hardcastle KA, Sethi D, Butchart A, Mikton C, et al. The effect of multiple adverse childhood experiences on health: a systematic review and meta-analysis. *Lancet Public Health*. 2017;2:e356–66.
- Mehta D, Kelly AB, Laurens KR, Haslam D, Williams KE, Walsh K, et al. Child maltreatment and long-term physical and mental health outcomes: an exploration of biopsychosocial determinants and implications for prevention. *Child Psychiatry Hum Dev*. 2021;54: 1–15.
- World Health Organization. Child maltreatment. 2024. Accessed September 5, 2024. <https://www.who.int/news-room/fact-sheets/detail/child-maltreatment#:~:text=Nonetheless%2C%20international%20studies%20reveal%20that,sexually%20abused%20as%20a%20child>
- Guajardo A, Tadros E. The long-term consequences of childhood maltreatment for adult survivors: a chronic price to pay. *J Psychol Perspect*. 2023;5:49–56.
- Felitti VJ, Anda RF, Nordenberg D, Williamson DF, Spitz AM, Edwards V, et al. Relationship of childhood abuse and household dysfunction to many of the leading causes of death in adults: the Adverse Childhood Experiences (ACE) Study. *Am J Prev Med*. 1998; 14:245–58.
- Brown DW, Anda RF, Tiemeier H, Felitti VJ, Edwards VJ, Croft JB, et al. Adverse childhood experiences and the risk of premature mortality. *Am J Prev Med*. 2009;37:389–96.
- Goodwin RD, Stein MB. Association between childhood trauma and physical disorders among adults in the United States. *Psychol Med*. 2004;34:509–20.
- Heim C, Binder EB. Current research trends in early life stress and depression: review of human studies on sensitive periods, gene-environment interactions, and epigenetics. *Exp Neurol*. 2012;233: 102–11.
- Nemeroff CB. Paradise lost: the neurobiological and clinical consequences of child abuse and neglect. *Neuron*. 2016;89:892–909.
- Norman RE, Byambaa M, De R, Butchart A, Scott J, Vos T. The long-term health consequences of child physical abuse, emotional abuse, and neglect: a systematic review and meta-analysis. *PLoS Med*. 2012;9:e1001349.
- Hammen C, Henry R, Daley SE. Depression and sensitization to stressors among young women as a function of childhood adversity. *J Consult Clin Psychol*. 2000;68:782.
- Shonkoff JP, Garner AS. Child CoPAo, Family Health CoEC, Adoption, Dependent Care, Developmental So, et al. The lifelong effects of early childhood adversity and toxic stress. *Pediatrics*. 2012; 129:e232–46.
- Valencia N, Seeger FR, Seitz KI, Carius L, Nkrumah RO, Schmitz M, et al. Childhood maltreatment and transdiagnostic connectivity of the default-mode network: the importance of duration of exposure. *J Psychiatr Res*. 2024;177:239–48.
- Elliot AJ, Mooney CJ, Infurna FJ, Chapman BP. Associations of lifetime trauma and chronic stress with C-reactive protein in adults ages 50 years and older: examining the moderating role of perceived control. *Psychosom Med*. 2017;79:622–30.
- Zhang H, Gertel VH, Cosgrove AL, Diaz MT. Age-related differences in resting-state and task-based network characteristics and cognition: a lifespan sample. *Neurobiol Aging*. 2021;101:262–72.
- Holz NE, Berhe O, Sacu S, Schwarz E, Tesarz J, Heim CM, et al. Early social adversity, altered brain functional connectivity, and mental health. *Biol Psychiatry*. 2023;93:430–41.
- Rebello K, Moura LM, Pinaya WH, Rohde LA, Sato JR. Default mode network maturation and environmental adversities during childhood. *Chronic Stress*. 2018;2:2470547018808295.
- Szeszko PR, Yehuda R. Magnetic resonance imaging predictors of psychotherapy treatment response in post-traumatic stress disorder: a role for the salience network. *Psychiatry Res*. 2019;277:52–7.
- Danese A, McEwen BS. Adverse childhood experiences, allostasis, allostatic load, and age-related disease. *Physiol Behav*. 2012;106:29–39.
- Glaser R, Kiecolt-Glaser JK. Stress-induced immune dysfunction: implications for health. *Nat Rev Immunol*. 2005;5:243–51.
- Baumeister D, Akhtar R, Ciufolini S, Pariante CM, Mondelli V. Childhood trauma and adulthood inflammation: a meta-analysis of peripheral C-reactive protein, interleukin-6 and tumour necrosis factor- $\alpha$ . *Mol Psychiatry*. 2016;21:642–9.
- Coelho R, Viola T, Walss-Bass C, Brietzke E, Grassi-Oliveira R. Childhood maltreatment and inflammatory markers: a systematic review. *Acta Psychiatr Scand*. 2014;129:180–92.
- Kerr DM, McDonald J, Minnis H. The association of child maltreatment and systemic inflammation in adulthood: a systematic review. *PLoS One*. 2021;16:e0243685.
- Danese A, Baldwin JR. Hidden wounds? Inflammatory links between childhood trauma and psychopathology. *Annu Rev Psychol*. 2017;68:517–44.
- Raison CL, Miller AH. Is depression an inflammatory disorder? *Curr Psychiatry Rep*. 2011;13:467–75.
- Busso DS, McLaughlin KA, Sheridan MA. Dimensions of adversity, physiological reactivity, and externalizing psychopathology in adolescence: deprivation and threat. *Psychosom Med*. 2017;79: 162–71.
- Infurna MR, Reichl C, Parzer P, Schimmenti A, Bifulco A, Kaess M. Associations between depression and specific childhood experiences of abuse and neglect: a meta-analysis. *J Affect Disord*. 2016;190: 47–55.
- Iob E, Lacey R, Steptoe A. The long-term association of adverse childhood experiences with C-reactive protein and hair cortisol: cumulative risk versus dimensions of adversity. *Brain Behav Immun*. 2020;87:318–28.
- Lacey R, Pereira SMP, Li L, Danese A. Adverse childhood

- experiences and adult inflammation: single adversity, cumulative risk and latent class approaches. *Brain Behav Immun.* 2020;87:820–30.
37. Cohen S, Wills TA. Stress, social support, and the buffering hypothesis. *Psychol Bull.* 1985;98:310.
  38. Brown S, Carbone JT, Hicks LM, Saini EK, Panisch LS, Dayton CJ. The moderating role of social support on the cortisol stress response of expectant fathers exposed to adverse childhood experiences. *J Fam Violence.* 2023;40:1–11.
  39. Helminen EC, Zhang X, Clawson AJ, Morton ML, Cary EL, Sinegar SE, et al. Stress-buffering effects of mindfulness programming for adolescents in schools during periods of high-and low-stress. *ECNU Rev Educ.* 2022;8:209–27.
  40. Bolton D. A revitalized biopsychosocial model: core theory, research paradigms, and clinical implications. *Psychol Med.* 2023;53:7504–11.
  41. Hobfoll SE. Social and psychological resources and adaptation. *Rev Gen Psychol.* 2002;6:307–24.
  42. Stinson DA, Fisher AN. Self-Esteem and Health. *The Wiley Encyclopedia of Health Psychology*, 1st edn. Wiley; 2020:615–21.
  43. Dogan T, Totan T, Sapmaz F. The role of self-esteem, psychological well-being, emotional self-efficacy, and affect balance on happiness: A path model. *Eur Sci J.* 2013;9:31–42.
  44. De Ridder D, Gillebaart M. Lessons learned from trait self-control in well-being: making the case for routines and initiation as important components of trait self-control. *Health Psychol Rev.* 2017;11:89–99.
  45. Carver CS, Scheier MF. Optimism, coping, and well-being. *The handbook of stress and health: A guide to research and practice*, 1st edn. Wiley; 2017: 400–14.
  46. Schiavon CC, Marchetti E, Gurgel LG, Busnelo FM, Reppold CT. Optimism and hope in chronic disease: a systematic review. *Front Psychol.* 2017;7:2022.
  47. Holt-Lunstad J, Smith TB, Baker M, Harris T, Stephenson D. Loneliness and social isolation as risk factors for mortality: a meta-analytic review. *Perspect Psychol Sci.* 2015;10:227–37.
  48. Uchino BN. Understanding the links between social support and physical health: a life-span perspective with emphasis on the separability of perceived and received support. *Perspect Psychol Sci.* 2009;4:236–55.
  49. Brewin CR, Andrews B, Valentine JD. Meta-analysis of risk factors for posttraumatic stress disorder in trauma-exposed adults. *J Consult Clin Psychol.* 2000;68:748.
  50. Ozer EJ, Best SR, Lipsey TL, Weiss DS. Predictors of posttraumatic stress disorder and symptoms in adults: a meta-analysis. *Psychol Bull.* 2003;129:52.
  51. Ciarrochi J, Morin AJ, Sahdra BK, Litalien D, Parker PD. A longitudinal person-centered perspective on youth social support: relations with psychological wellbeing. *Dev Psychol.* 2017;53:1154.
  52. Slemp GR, Field JG, Ryan RM, Forner VW, Van den Broeck A, Lewis KJ. Interpersonal supports for basic psychological needs and their relations with motivation, well-being, and performance: a meta-analysis. *J Pers Soc Psychol.* 2024;127:1012–37.
  53. Sperry DM, Widom CS. Child abuse and neglect, social support, and psychopathology in adulthood: a prospective investigation. *Child Abuse Neglect.* 2013;37:415–25.
  54. Finkelhor D, Turner H, Ormrod R, Hamby SL. Violence, abuse, and crime exposure in a national sample of children and youth. *Pediatrics.* 2009;124:1411–23.
  55. Fisher HL, Caspi A, Moffitt TE, Wertz J, Gray R, Newbury J, et al. Measuring adolescents' exposure to victimization: the environmental risk (E-Risk) longitudinal twin study. *Dev Psychopathol.* 2015;27(pt 2):1399–416.
  56. Salazar AM, Keller TE, Courtney ME. Understanding social support's role in the relationship between maltreatment and depression in youth with foster care experience. *Child Maltreat.* 2011;16:102–13.
  57. Horan JM, Widom CS. From childhood maltreatment to allostatic load in adulthood: the role of social support. *Child Maltreat.* 2015; 20:229–39.
  58. Ryff CD, Seeman T, Weinstein M. *Midlife in the United States (MIDUS 2): Biomarker Project*. Inter-university Consortium for Political and Social Research [distributor]; 2022.
  59. Love GD, Seeman TE, Weinstein M, Ryff CD. Bioindicators in the MIDUS national study: protocol, measures, sample, and comparative context. *J Aging Health.* 2010;22:1059–80.
  60. Bernstein DP, Stein JA, Newcomb MD, Walker E, Pogge D, Ahluvalia T, et al. Development and validation of a brief screening version of the Childhood Trauma Questionnaire. *Child Abuse Neglect.* 2003;27:169–90.
  61. Burns C, Hejl C, Z Szabo Szabo Y. Childhood adversity and adult inflammation: exploring the mediating role of emotion regulation in the MIDUS II study. *J Child Adolesc Trauma.* 2024;17:319–34.
  62. McLaughlin KA, Sheridan MA. Beyond cumulative risk: a dimensional approach to childhood adversity. *Curr Dir Psychol Sci.* 2016; 25:239–45.
  63. Lachman ME, Weaver SL. The sense of control as a moderator of social class differences in health and well-being. *J Pers Soc Psychol.* 1998;74:763.
  64. Rosenberg M. Rosenberg self-esteem scale (RSE). *Acceptance and commitment therapy. Measures package.* 1965;61:18.
  65. Scheier MF, Carver CS. Effects of optimism on psychological and physical well-being: theoretical overview and empirical update. *Cognit Ther Res.* 1992;16:201–28.
  66. Stokes JE. Social integration, perceived discrimination, and self-esteem in mid- and later life: intersections with age and neuroticism. *Aging Mental Health.* 2019;23:727–35.
  67. Rook KS. Parallels in the study of social support and social strain. *J Soc Clin Psychol.* 1990;9:118–32.
  68. Radloff LS. The CES-D scale: a self-report depression scale for research in the general population. *Appl Psychol Measure.* 1977;1: 385–401.
  69. Ryff CD. Happiness is everything, or is it? Explorations on the meaning of psychological well-being. *J Pers Soc Psychol.* 1989;57: 1069.
  70. Boylan JM, Ryff CD. Psychological well-being and metabolic syndrome: findings from the Midlife in the United States national sample. *Psychosom Med.* 2015;77:548–58.
  71. Friedman EM, Herd P. Income, education, and inflammation: differential associations in a national probability sample (The MIDUS study). *Psychosom Med.* 2010;72:290–300.
  72. Austin AE, Lesak AM, Shanahan ME. Risk and protective factors for child maltreatment: a review. *Curr Epidemiol Rep.* 2020; 7:334–42.
  73. Muthén B, Muthén L. *Mplus User's Guide*, 8th edn. Los Angeles, CA: Muthén & Muthén; 1998-2017.
  74. Hu LT, Bentler PM. Evaluating Model Fit. *Structural Equation Modelling Concepts, Issues, and Applications*, 1st. Thousand Oaks, CA: Sage; 1995: 76–99.
  75. Bermudez EA, Rifai N, Buring J, Manson JE, Ridker PM. Interrelationships among circulating interleukin-6, C-reactive protein, and traditional cardiovascular risk factors in women. *Arterioscler, Thromb, Vasc Biol.* 2002;22:1668–73.
  76. Enders CK, Bandalos DL. The relative performance of full information maximum likelihood estimation for missing data in structural equation models. *Struct Equ Model.* 2001;8:430–57.
  77. Renna ME, Peng J, Shrout MR, Madison AA, Andridge R, Alfano CM, et al. Childhood abuse histories predict steeper inflammatory trajectories across time. *Brain Behav Immun.* 2021;91:541–5.
  78. Gilbert P, McEwan K, Mitra R, Franks L, Richter A, Rockliff H. Feeling safe and content: a specific affect regulation system? Relationship to depression, anxiety, stress, and self-criticism. *J Positive Psychol.* 2008;3:182–91.
  79. Cicchetti D, Toth SL. Child maltreatment and developmental psychopathology: a multilevel perspective. *Dev Psychopathol.* 2016; 3:457–512.
  80. Teicher MH, Gordon JB, Nemeroff CB. Recognizing the importance of childhood maltreatment as a critical factor in psychiatric diagnoses, treatment, research, prevention, and education. *Mol Psychiatry.* 2022;27:1331–8.
  81. Hill TD, Kaplan LM, French MT, Johnson RJ. Victimization in early life and mental health in adulthood: an examination of the mediating and moderating influences of psychosocial resources. *J Health Soc Behav.* 2010;51:48–63.
  82. Lee ST, Choy BK, Yong JC. A bubble of protection: examining

- dispositional optimism as a psychological buffer of the deleterious association between negative work-family spillover and psychological health. *Eur J Work Organ Psychol.* 2022;31:35–46.
83. Magin ZE, Park CL, Burke JD, Infurna FJ. Perceived control and inflammation: mediating and moderating effects in the relationship between cumulative trauma and depression. *Psychosom Med.* 2024;86:192–201.
84. Lagdon S, Ross J, Robinson M, Contractor AA, Charak R, Armour C. Assessing the mediating role of social support in childhood maltreatment and psychopathology among college students in Northern Ireland. *J Interpers Violence.* 2021;36:NP2112–36.
85. Struck N, Krug A, Feldmann M, Yuksel D, Stein F, Schmitt S, et al. Attachment and social support mediate the association between childhood maltreatment and depressive symptoms. *J Affect Disord.* 2020;273:310–7.
86. Fagundes CP, Glaser R, Kiecolt-Glaser JK. Stressful early life experiences and immune dysregulation across the lifespan. *Brain Behav Immun.* 2013;27:8–12.
87. Arslan G. Psychological maltreatment and substance use among college students: psychological distress, belongingness, and social support. *J Ethn Subst Abuse.* 2024;23:426–49.
88. Baldwin JR, Reuben A, Newbury JB, Danese A. Agreement between prospective and retrospective measures of childhood maltreatment: a systematic review and meta-analysis. *JAMA Psychiatry.* 2019;76:584–93.
89. Newbury JB, Arseneault L, Moffitt TE, Caspi A, Danese A, Baldwin JR, et al. Measuring childhood maltreatment to predict early-adult psychopathology: comparison of prospective informant-reports and retrospective self-reports. *J Psychiatr Res.* 2018;96:57–64.
90. Shenk CE, Shores KA, Ram N, Felt JM, Chimed-Ochir U, Olson AE, et al. Contamination in observational research on child maltreatment: a conceptual and empirical review with implications for future research. *Child Maltreat.* 2025;30:9–20.
91. Nouman H, Alfandari R, Enosh G, Dolev L, Daskal-Weichhendler H. Mandatory reporting between legal requirements and personal interpretations: community healthcare professionals' reporting of child maltreatment. *Child Abuse Neglect.* 2020;101:104261.